natural plant nutrient supplement

CROPMAX is a growth stimulant for all kinds of crops. The activity is based on the combination of trace elements, amino acids, vitamins and polysaccharides.

CROPMAX stimulates growth in different stages of plantfysiology:

- 1. At germination of seed and youngplants, promoting growth during the vulnerable early stages of growth
- 2. At root setting (forming a better root system and thus improving feed and water uptake)
- 3. During the vegetative growth when this is required (lettuce, spinach, tobacco, roses, tea, etc.)
- 4. During the generative stages, increasing the amount of flowers and promoting fruitsetting
- 5. During maturation, promoting formation of dry matter (and oil in oilseeds, etc.)



Ananas (Ananas)

Make a solution of 1 ltr. with a sufficient amount of water to spray 1 hectare (between 500 and 1000 ltr. depending on the sprayer capacity).

- Apply when the plants set roots at the earlier stage of vegetation.
- Apply a second time when the plants enter the generative stadium (at flower-set)
- A third application needs to be done 4 weeks after the second application.







The dilution rate of **CROPMAX** with water depends on the calibration of the sprayer. It can vary between 100 and 500 litre per hectare. This depends on the pressure of the sprayer, the size of the spraying nozzle and the speed of the sprayer across the treated area. The calibration of the sprayer can be thus influenced, depending on the density of treated crops (high planting density, more solution).

NOTE: DO NOT SPRAY, when the sun is shining intensely, because than the plant closes the pores, making the uptake of active ingredient less efficient. Therefore, spray on sunny days in the morning or at dusk. The time used for taking up the active ingredient is 4 hours max.